

## Investigation of toxic effect of *moringa oleifera* leaf-methanol extract on breast cancer cell mcf-7

Nazia Hossain, Mohamed Elwathig Saeed Mirghani and Raha Raus

Nazia Hossain, B.Eng of Biochemical-Biotechnology Engineering, Faculty of Biochemical-Biotechnology Engineering, International Islamic University Malaysia, Malaysia.

Associate Professor Dr. Mohamed Elwathig Saeed Mirghani, Department of Biotechnology Engineering, Kulliyah (Faculty) of Engineering, International Islamic University Malaysia, 53100, Jalan Gombak, Malaysia. International Institute for Halal Research and Training (INHART), IIUM, Gombak, Malaysia

Assistant Professor Raha Bt. Raus, Department of Biotechnology Engineering, Kulliyah (Faculty) of Engineering, International Islamic University Malaysia, 53100, Jalan Gombak, Malaysia.

### Abstract

*Moringa oleifera* (*M. oleifera*) leaf extracted with methanol was screened for anti-breast cancer activity. Microtiter tetrazolium (MTT) assay was used to determine the anti-cancer activity. MTT assay was evaluated on two types of cell lines normal healthy cell-line VERO and breast cancer cell-line MCF-7. Both cell-lines were seeded at  $2 \times 10^5$  cells per milliliter (ml) in a 100 $\mu$ l of media (Dulbecco Modified Eagle Medium 90% and serum 10%) per well in 96 well plates incorporation with 4 $\mu$ l/ml *M. oleifera* leaf extract at wavelength  $A_{570nm}$  and 37 $^{\circ}$  C in CO<sub>2</sub> incubator (with 5% CO<sub>2</sub> and 95% air) which is the physiological temperature. The highest cancer cell inhibition rate was 88.39% ( $1.77 \times 10^5$  cells per ml). Moreover, *M.oleifera* leaf extraction conditions were optimized in terms of temperature and incubation time by sonication method using Design of Experiment (DoE) software with response surface methodology. The optimum condition for leaf extraction was measured to be temperature 50 $^{\circ}$ C and incubation time 45 min with medium frequency and it obtained 87.13% ( $1.74 \times 10^5$  cells per ml) cancer cell growth inhibition and 13.92% ( $27.8 \times 10^3$  cells per ml) normal cell inhibition by this condition.

**KEYWORDS:** *Moringa medicinal potential, Anti-Cancer Activity, Breast Cancer Inhibition, Optimization of Moringa leaf extract, Anti-Breast Cancer Drug Discovery, Toxic Effect of Moringa leaf against MCF-7*

### Corresponding Author:

Nazia Hossain, B.Eng of Biochemical-Biotechnology Engineering  
Faculty of Biochemical-Biotechnology Engineering, International Islamic  
University Malaysia, Malaysia.  
Email: [bristy808.nh@gmail.com](mailto:bristy808.nh@gmail.com) & [bristy808@yahoo.com](mailto:bristy808@yahoo.com)

## INTRODUCTION

Breast cancer is the most leading cause of women mortality in whole world. 7.6 million people are dying worldwide by cancer every year where 1.38 million of breast cancer cases, the highest death rate cause. The World Health Organization (WHO) of the United Nations (UN) projects that without immediate action, the global number of deaths from cancer will increase by nearly 80% by 2030, with most occurring in low and middle income country. (Abeloff, 2008) & (GLOBOCAN, 2014)

DNA carries genes that are the blueprints for protein produced in the body and proteins work to suppress the growth. But sometimes mutation occurs in some particular genes because of radiation, Ultra-violet radiation, smoking, alcohol or other carcinogens what can cause the abnormal production of proteins and the combination of these abnormal proteins create cancer as a result (Blask, 1988). Cancer cells flock and build up tumors that may compress, invade and cause destruction of normal tissues. Sometimes, cells break away from the tumor and flow through the bloodstream or the lymph system to others organs of body parts and settle down somewhere and stay permanently to build up 'colony tumor' and the cells start to grow over there. This new site of tumor forming is known as 'Metastasis' (American Society of Clinical Oncology, 2009) (Shishodia, 2004).

Plant-derived substances have recently achieved great interest owing to their versatile applications such as *Moringa oleifera* of monogenic family, the *Moringaceae*. It is a well famous plant cultivated in tropical or sub-tropical areas such as Himalayan tracts of India, Pakistan, Bangladesh, Afganistan, Malaysia, Indonesia, Egypt, Sudan, Nigeria, Chad, Somalia, Tanzania and others. It is known as 'drumstick tree', 'golden shower tree', 'horseradish tree', 'ben oil tree' and 'benzoin tree' (Musa, 2012). The leaves contain nutrients especially essential amino acids, vitamins,

minerals,  $\beta$ -carotene, benzyl isothiocyanate, niazimicin, pterygospermin, benzyl isothiocyanate, benzyl glucosinolate. Besides nutritional benefits, *M. oleifera* is used for the treatment of rheumatism, ascites, infection, hiccough influenza and internal abscess (Anwar, 2007).

Few researches were gone through different types of cancer cells incorporation with ethanol extraction of *Moringa* leaves and they showed strong cytotoxicity against cancer cells such as hepatocarcinoma and coloractal adenocarcinoma. That's why *Moringa* leaves extraction was experimented to inhibit breast cancer cells (MCF-7) in this study (Hossain & et al., 2015) (Charoensin, 2014). Compare to ethanol extraction, methanol extraction is easier, less time consuming. Methanol was normally preferred in this study as the organic solvent since it is more polar as compared to other types of alcohol such as ethanol, acetone etc and in many cases methanol extraction produced the best extraction results. Especially to extract leaves in different conditions by sonication (cell disruption by ultra-sound), only methanol showed stable extraction where ethanol and acetone caused cover-bursting effect with high temperature (King, 1984).

To mention, differents parts of *M. oleifera* have been experimented to investigate its' effect on various types of diseases. Nevertheless, there is limited research for *M. oleifera* leaf extract on anti breast cancer cell and there is no methanol leaf extraction on breast cancer cell has been experimented. So, the goal of this research was to investigate anti-breast cancer activity of *Moringa* leaves with methanol extraction by sonication method and optimize it with different conditions run by Design of Experiment (DoE) optimization software (Box, 2010) (Charoensin, 2014).

## MATERIALS AND METHODS

**Materials:** Fresh *Moringa oleifera* leaves were collected from a garden near to Terminal Putra LRT station, Gombak, Kuala Lumpur, Malaysia. Experimented samples both cell-lines breast cancer cell-line (MCF-7) & normal cell-line (VERO) and the chemicals accutase, phosphate buffer solution (PBS) 10x solution, DMEM (Dulbecco Modified Eagle Medium) powder, 70% ethanol for sterilization, Fetal Bovine Serum (FBS), dimethyl sulphoxide (DMSO), NaOH powder or pellet, HCl, Sodium carbonate powder, tryptophan blue dye & MTT assay were collected from IIUM cell & tissue laboratory.

**Methods:** Fresh *Moringa oleifera* leaves were washed with clean water, dried, ground and stored for experiment. The *Moringa* leaf-methanol extraction process was optimized with 11 runs done by Design Expert software 6.0.8 (DOE) using response surface methodology. Extraction (11 runs) of the leaves was done by sonication method through an ultrasonic bath at medium frequency at 40-60°C for 30-60 minutes. 3g of each dried sample was dissolved in 30ml of methanol. The crude extract was isolated through vacuum filtration and rotary evaporation process. Then extract was transferred to a 15ml tube and 10ml DMSO was added to preserve. Normal cell-line (VERO) and Breast cancer cell-line (MCF-7) were cultured through stored frozen cells thawing, media change, sub-culturing monolayer cells and cell counting respectively. The cells were seeded at  $2 \times 10^5$  cells per ml for breast cancer cell in a 100 $\mu$ l of media (DMEM 90% and serum 10%) per well in 96 well plates. Both cell-lines were incorporated with 4 $\mu$ l/ml *Moringa* leaves-methanol extract in different plates and finally MTT assay was added on the incorporation and the absorbance at A570 nm was recorded immediately by an ELISA plate reader. Then the data was analyzed.

## RESULTS

### Optimization of Process Conditions by Design of Expert (DOE)

To optimize the cancer cell inhibition it in terms of temperature and incubation time. 11 experiments were run as designed using Design of Expert (DOE) software, version 6.0.8 to attain its optimal yield at 50°C, at an optimal incubation time of 45 minute.

### Anova Analysis

#### Response: Cell growth Inhibition

Analysis of Variance (ANOVA) for Response Surface Quadratic Model was used to determine whether the model is significant or not. The coefficient p-value for linear, quadratic and interaction terms are shown in Table 1.

The Model F-value of 22.79 implies the model is significant. There is only a 0.19% chance that a "Model F-Value" this large could occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case B, A<sup>2</sup>, B<sup>2</sup> are significant model terms. Values greater than 0.1000 indicate the model terms are not significant. The "Lack of Fit F-value" of 1.12 implies the Lack of Fit is not significant relative to the pure error. There is a 50.42% chance that a "Lack of Fit F-value" this large could occur due to noise.

The "Pred R-Squared" of 0.7142 is not as close to the "Adj R-Squared" of 0.9159 that was very much expected. A ratio greater than 4 is desirable (Box, 2010). Here, the ratio of 13.743 indicates an adequate signal for successful optimized design. Mean is the average value of all outcomes and here mean cancer cell growth inhibition percentage by *Moringa oleifera* leaves extract is 87.13 what is very positive with the project.

$$C. V. \% = \frac{Std. Dev}{Mean} \times 100\%$$

**Table 1:** Sequential Model Sum of Squares

Source	Sum of Squares	df	Mean Square	F value	P value Prob > F	
<b>Model</b>	163.58	5	32.72	22.79	0.0019	Significant
<b>A-Temperature</b>	001.68	1	01.68	01.17	0.3289	
<b>B-Incubation</b>	088.26	1	88.26	61.49	0.0005	Significant
<b>AB</b>	003.70	1	03.70	02.58	0.1692	
<b>A<sup>2</sup></b>	019.11	1	19.11	13.31	0.0148	Significant
<b>B<sup>2</sup></b>	064.59	1	19.11	45.00	0.0011	Significant
<b>Residual</b>	007.18	5	64.59			
<b>Lack of fit</b>	004.50	3	01.50	01.12	0.5042	Not Significant
<b>Pure Error</b>	002.68	2	01.34			
<b>Core Total</b>	170.76	10				

**Table 2:** Model characteristics

<b>Std. Dev.</b>	1.20	<b>R-Squared</b>	0.9580
<b>Mean</b>	87.13	<b>Adj R-Squared</b>	0.9159
<b>C.V.%</b>	1.37	<b>Pred R-Squared</b>	0.7142
<b>PRESS</b>	48.81	<b>Adeq Precision</b>	13.743

**Table 3:** VERO cell-line growth Inhibition by different conditions

Temperature (°C)	Incubation Time (min)	Cell Growth Inhibition (%)
50	45	19.40
50	30	6.08
40	60	12.17
50	45	13.92

In this project, C.V. % was very low 1.37 what good sign for result stability. The predicted residual sum of squares (PRESS) was 48.81 that was less than 50.

**Diagnostic**

In the diagnostic, Figure 1 (predicted vs. actual response) showed that the distribution of points are closed to linear line (R<sup>2</sup> = 0.9580) and Figure 2 shows the 3 dimensional view of the whole optimization process and optimized point as well.

**Experiment Result**

The results were taken using ELISA microplate reader at 570nm. In order to validate the result obtained from triplicate for each run. Thus using the same concentration range, three 96 well plates were done for each run of each cell-line. Average was taken to obtain the final value and cell viability and cell growth inhibition percentage for both types of cells were calculated by the equations:

$$\text{Growth} = \frac{OD_{\text{treated cells}}}{OD_{\text{untreated cells}}} \times 100\%$$

$$\text{Growth inhibition} = 100\% - \frac{OD_{\text{treated cells}}}{OD_{\text{untreated cells}}} \times 100\%$$

OD *untreated cells* = Media with cells without extract

OD *treated cells* = Media with cells with extract

For normal cell-line (VERO), four experiments were done with different conditions based on DoE table to check *M. oleifera* leaves extract affect normal cells or not. These all experiments showed normal cell inhibition by average cell inhibition 12.89% and in optimum extraction condition at temperature 50°C and incubation time 45 min with medium frequency, cell growth inhibition is 13.92% (27.8 x 10<sup>3</sup> cells per ml).

There were 11 runs of experiments done on MCF-7 by *Moringa* leaves extract with different conditions based on time and temperature in terms of DOE table and analyzed the results. It is shown in Table 4..

Table 4 has shown high cancer cell growth inhibition by the extract. Average range of growth inhibition has been 80% - 90% (mean value 87.13%) with various conditions and in optimum extraction condition at temperature 50°C and incubation time 45 min with medium frequency, cell growth inhibition is 88.39% (1.77 x 10<sup>5</sup> cells per ml).




Figure 1: Predicted vs. Actual values




Figure 2: 3D view plot for desirability

Table 4: Result Analysis by DOE

Std	Run	Factor 1 A : Temperature (°C)	Factor 2 B : Incubation Time (min)	Response 1 Cell Growth Inhibition (%)
3	1	50	60	78.4659
4	2	60	60	80.8468
2	3	60	30	89.7957
1	4	40	30	89.7979
6	5	60	45	91.0828
9	6	50	45	87.3738
5	7	40	45	90.4039
10	8	50	45	89.6298
7	9	50	30	87.4286
3	10	40	60	84.6971
11	11	50	45	88.9549

## DISCUSSION

Optimization is a process to find an alternative with the most cost-effective or highest achievable performance under the given constraints by maximizing desired factors and minimizing undesired ones (Hossain & et al., 2015) (Porter & et al., 1997). To optimize this extraction process, DoE statistical analyses was used in this study where average PRESS was 48.81. Average PRESS value is the determination factor whether the model is over-fitted or under-fitted and PRESS value less than 50 is desired for optimized design (Box, 2010). So, in this study optimized design was good overall. Besides, predicted vs. actual response showed that the distribution of points are closed to linear line ( $R^2 = 0.9580$ ). As difference between the actual and predicted value is small, the result is counted better (Box, 2010). In this design, difference for cell growth inhibition is very small in all runs that indicated the model is very good. These results proved that Response Surface Methodology statistical analysis fitted this experiment for optimizing the extraction condition.

*Moringa oleifera* extract work on different cancer cells in other research (Charoensin, 2014). In this research, breast cancer cell (MCF-7) growth inhibition was 87.13% ( $1.74 \times 10^5$  cells per ml) in average that proved *Moringa* leaves extract was able to inhibit breast cancer cell with very high percentage in all kinds of condition parameters. When mixture was diluted with media the growth inhibition was also reduced, which evidenced that *Moringa* leaves extract was very strong against MCF-7 cell growth.

On the other hand, in this research, experiments on normal cell-lines (VERO) showed that *Moringa* extract inhibited VERO cell-line with small percentage too. But based on previous researches, it was proven that *Moringa oleifera* is good for health nutrition and its leaves extract usually does not kill normal cell-line (Anwar, 2007). In this case, it was suspected that the presence of methanol with leaves extract might affect this result by inhibiting normal cell growth. Because methanol is volatile alcohol similar with drinking alcohol and sometimes

methanol causes toxicity that may cause metabolic acidosis, neurologic squealed and even death if ingested (Mukherjee, 2001). To avoid this impurity, DMSO is recommended to use as solvent to re-dissolve the extract instead of methanol though using DMSO is expensive. For further research, dose responsive experiment should be recommended.

The MTT assay used to evaluate the cytotoxicity of methanolic leaf extract of *Moringa oleifera* on MCF-7 and VERO cell lines showed that the extract is significantly cytotoxic to MCF-7 cell line and considerably less so towards normal cell-line VERO. These findings highlights the potentials of *Moringa oleifera* leaf extract in the treatment of breast cancer. Its therapeutic potential is huge and can be used as alternative to or supplementation for the various therapies. However there is a need to identify the actual components responsible for this cytotoxicity and to isolate them and to study their effect in vivo to ascertain their efficacies and or any side effects. The concerns of side effects with pharmacological agents, and growing interest in plant bio-resources for treatment of cancers mean this extract could have important role in future studies on the management of breast cancer.

## CONCLUSION

The *Moringa* leaf-methanol extract showed extremely high anti-breast cancer activity that indicates the medicinal value of *M. oleifera* in terms of cancer preventive drugs and chemotherapy. The *Moringa* leaves extraction optimum condition was found to be temperature  $50^\circ\text{C}$  and incubation time 45 min with medium frequency at sonication. It is strongly suggested that it could potentially be an ideal anticancer therapeutic candidate specific to cancer cells. These results also suggested that extraction strategy adopted in this work could lead to several industrial applications and the potential therapeutic implications of the soluble extract from *M. oleifera* leaves extract in the treatment of various types of cancers.

## ACKNOWLEDGEMENT

This work supported by the Cell and Tissue Laboratory at Biochemical-Biotechnology Engineering Department (BTE), Faculty of Engineering, International Islamic University Malaysia (IIUM)

## REFERENCES

- Abeloff MD, Wolff AC and Weber BL (2008). Cancer of the Breast. *Clinical Oncology*. 4th ed. Philadelphia, Elsevier, 1875–1943.
- American Society of Clinical Oncology. Clinical Cancer Advances 2009. Major Research Advances in Cancer Treatment, Prevention and Screening
- Blask, S. M. (1988). Effects of the Pineal Hormone Melatonin on the Proliferation and Morphological Characteristics of Human Breast Cancer Cells (MCF-7) in Culture. *Cancer Research* , **48**, 6121-6126.
- Box, G. a (2010). On the Experiment Attainment of Optimum Conditions. *Journal of the royal Statistical Society Series B* , **13** (1), 1-45.
- Charoensin, S. (2014).Antioxidant and anticancer activities of *Moringa oleifera* leaves. *Journal of Medicinal Plant Research* , **8**(7), pp 318-325
- Farooq Anwar, S. L. (2007). Moringa oleifera: A Food Plant with Multiple Medicinal Uses. *PHYTOTHERAPY RESEARCH* , **21**, 17-25.
- GLOBOCAN (2014). Breast Cancer Incidence and Mortality Worldwide, Cancer Incidence in Five Continents (C15) Volume I-X, International Agency for Research on Cancer, World Health Organization.
- King, C. L. (1984). Factor Influencing Solvent Selection for Extraction of Methanol and Ethanol from Aqueous Solution. *Ind.Eng.Chemical Process Des.* , **1** (23), 109-115.
- Mukherjee, A. (2001). *Advances in Cancer therapy with Plant based Natural Products* , **8**, 1467-1486.
- Musa, E. N. (2012). An investigation of Anti-cancer activity of *Moringa oleifera* leaves. 1-5.
- Hossain, N., Mirghani, M., & Raus, R. (2015). Optimization of *Moringa oleifera* Leaf Extraction and Investigation of Anti Breast Cancer Activity with the Leaf Extract. *Engineering International* , **3**(2). Retrieved from <http://www.journals.abc.us.org/index.php/ei/article/view/775>
- Porter S.C, Verseput R.P. and Cunnigham C.R. (October, 1997). "Process Optimization using Design of Experiments". *Pharm. Technology* **1**.
- Shishodia, S. a. (2004). Guggulsteron Inhibits nf-kb and ikbs-kinase activation. *Suppresses Expression of Anti-Apototic Gene Products and Enhances Apoptosis* , 279.

Supplementary Data




Figure 1: Contour view plot for desirability




Figure 2: 3D view plot for desirability